

φX174 RF II DNA



1-800-632-7799
info@neb.com
www.neb.com



N3022L 124160418041

N3022L

150 µg **Lot: 1241604** **Exp: 4/18**

1,000 µg/ml **Store at -20°C**

Description: This is the double-stranded, nicked, circular form of φX174 DNA (relaxed form). The molecular weight of φX174 DNA is 3.50 X 10⁶ daltons it is 5,386 base pairs in length. Greater than 90% of the molecules are RF II, the remainder are RF I (supercoiled) and linear φX174 DNA.

Supplied in: 10 mM Tris-HCl (pH 8.0), 1 mM EDTA.

Preparation: φX174 RF II is prepared by relaxing (nicking) φX174 RF I DNA in high salt at 75°C. Greater than 90% of the molecules are in the RF II form as indicated by agarose gel electrophoresis.

References:

1. Sanger, F. et al. (1978) *J. Mol. Biol.* 125, 225.
2. Freifelder, D. and Dewitt, R. (1977) *Gene* 1, 385.

CERTIFICATE OF ANALYSIS

φX174 RF II DNA



N3022L 124160418041

N3022L

150 µg **Lot: 1241604** **Exp: 4/18**

1,000 µg/ml **Store at -20°C**

Description: This is the double-stranded, nicked, circular form of φX174 DNA (relaxed form). The molecular weight of φX174 DNA is 3.50 X 10⁶ daltons it is 5,386 base pairs in length. Greater than 90% of the molecules are RF II, the remainder are RF I (supercoiled) and linear φX174 DNA.

Supplied in: 10 mM Tris-HCl (pH 8.0), 1 mM EDTA.

Preparation: φX174 RF II is prepared by relaxing (nicking) φX174 RF I DNA in high salt at 75°C. Greater than 90% of the molecules are in the RF II form as indicated by agarose gel electrophoresis.

References:

1. Sanger, F. et al. (1978) *J. Mol. Biol.* 125, 225.
2. Freifelder, D. and Dewitt, R. (1977) *Gene* 1, 385.

CERTIFICATE OF ANALYSIS