

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

LAMP Fluorescent Dye
B1700S
50 X Concentrate
10159436
11/2023
-20°C
PS-B1700S v1.0
Proprietary

LAMP Fluorescent Dye Component List					
NEB Part Number	Component Description	Lot Number	Individual QC Result		
B1700SVIAL	LAMP Fluorescent Dye	10135817	Pass		

Assay Name/Specification	Lot # 10159436
Functional Testing (LAMP, Master Mix) A 25 μ I reaction with 1X WarmStart® LAMP Master Mix in the presence of 1X LAMP Primers containing 10 ng genomic DNA and 1X LAMP fluorescent dye results in a threshold time of ≤ 20 minutes as determined by fluorescent detection.	Pass
Endonuclease Activity (Nicking) A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 2 μ l of LAMP Fluorescent Dye incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2 µl of LAMP Fluorescent Dye incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of LAMP Fluorescent Dye is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Functional Testing (RT-LAMP, Master Mix)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10159436
A 25 μ I reaction with 1X WarmStart® LAMP Master Mix in the presence of 1X LAMP Primers containing 10 ng of genomic RNA and 1X LAMP fluorescent dye results in a threshold time of ≤ 20 minutes as determined by fluorescent detection.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 μ I of LAMP Fluorescent Dye is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is \leq 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Vanguez histi

Christie Vazquez Production Scientist 20 Jul 2022

Erin Varney

Packaging Quality Control Inspector 20 Jul 2022

