

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | NEBuffer™ 2.1 |
|------------------------|--|
| Catalog Number: | B7202S |
| Concentration: | 10 X Concentrate |
| Packaging Lot Number: | 10070033 |
| Expiration Date: | 02/2023 |
| Storage Temperature: | -20°C |
| Specification Version: | PS-B7202S v1.0 |
| Composition (1X): | 50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 100 μg/ml BSA, (pH 7.9 @ 25°C) |

| NEBuffer™ 2.1 Component List | | | | |
|------------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| B7202SVIAL | NEBuffer™ 2.1 | 10067776 | Pass | |

| Assay Name/Specification | Lot # 10070033 |
|---|----------------|
| RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 2.1 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection. | Pass |
| pH (buffers/solutions) The pH of 10X NEBuffer 2.1 is between pH 7.8 and 8.0 at 25°C. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10070033 |
|--|----------------|
| Endonuclease Activity (Nicking, Buffer) A 50 μ I reaction in 1X NEBuffer 2.1 containing 1 μ g of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Conductivity (buffers/solutions) The conductivity of 10X NEBuffer 2.1 is between 55 and 62 mS at 25°C. | Pass |

This product has been tested and shown to be in compliance with all specifications.

Production Scientist 14 Feb 2020

ray-

An Minichiello Packaging Quality Control Inspector 19 Mar 2020

