

www.neb.com info@neb.com



New England Biolabs Certificate of Analysis

Product Name: DNA Polymerase I, Large (Klenow) Fragment

Catalog Number: M0210M
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10

nmol of dNTP into acid insoluble material in 30 minutes at 37°C.

Packaging Lot Number: 10221782
Expiration Date: 12/2025
Storage Temperature: -20°C

Storage Conditions: 25 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @

25°C)

Specification Version: PS-M0210M v1.0

DNA Polymerase I, Large (Klenow) Fragment Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0210MVIAL	DNA Polymerase I, Large (Klenow) Fragment	10221783	Pass	
B7002SVIAL	NEBuffer™ 2	10221172	Pass	

Assay Name/Specification	Lot # 10221782
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of DNA Polymerase I, Large (Klenow) Fragment incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units DNA Polymerase I, Large (Klenow) Fragment incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) DNA Polymerase I, Large (Klenow) Fragment is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of DNA Polymerase I, Large (Klenow) Fragment is incubated at	Pass



M0210M / Lot: 10221782

Page 1 of 2



Assay Name/Specification	Lot # 10221782
37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 50 units of DNA Polymerase I, Large (Klenow) Fragment is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lea Antonopoulos

Production Scientist

31 Jan 2024

Michael Tonello

Packaging Quality Control Inspector

02 Apr 2024



M0210M / Lot: 10221782

Page 2 of 2