

New England Biolabs Certificate of Analysis

Product Name: Quick-Load[®] Taq 2X Master Mix
Catalog #: M0271S/L
Concentration: 2X Concentrate
Lot #: 0371707
Assay Date: 07/2017
Expiration Date: 07/2019
Storage Temp: -20°C
Composition (1X): 10 mM Tris-HCl (pH 8.6 @ 25°C), 50 mM KCl, 1.5 mM MgCl₂, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 0.08 % IGEPAL[®] CA-630, 0.05 % Tween[®] 20, 0.024 % Orange G, 0.0025 % Xylene cyanol, 33 units/ml Taq DNA Polymerase
Specification Version: PS-M0271S/L v1.0
Effective Date: 10 May 2017

| Assay Name/Specification (minimum release criteria) | Lot #0371707 |
|--|--------------|
| Endonuclease Activity (Nicking) - A 50 µl reaction in ThermoPol [®] Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at either 37°C or 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 1X Quick-Load [®] Taq Master Mix containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| PCR Amplification (5 kb Lambda, Master Mix) - A 25 µl reaction in 1X Quick-Load [®] Taq Master Mix and 0.2 µM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification results in the expected 5 kb product. | Pass |
| Phosphatase Activity (pNPP) - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Taq DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis. | Pass |
| Protein Purity Assay (SDS-PAGE) - Taq DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |



New England Biolabs Certificate of Analysis

| Assay Name/Specification (minimum release criteria) | Lot #0371707 |
|--|--------------|
| <p>qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 5 units of <i>Taq</i> DNA Polymerase is screened for the presence of <i>E. coli</i> genomic DNA using SYBR[®] Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.</p> | Pass |
| <p>RNase Activity (Extended Digestion) - A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of Quick-Load[®] <i>Taq</i> 2X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | Pass |
| <p>Single Stranded DNase Activity (FAM-Labeled Oligo) - A 50 μl reaction in ThermoPol[®] Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of <i>Taq</i> DNA Polymerase incubated for 30 minutes at either 37°C or 75°C yields <10% degradation as determined by capillary electrophoresis.</p> | Pass |



Authorized by
Karen Moreira
10 May 2017



Inspected by
Tony Spear-Alfonso
01 Aug 2017

