

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: RNase H
Catalog Number: M0297S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to produce 1

nmol of ribonucleotides from 20 picomoles of a fluorescently labeled 50 base pair RNA-DNA hybrid in a total reaction volume of 50 μ l in

20 minutes at 37°C.

Packaging Lot Number: 10149191
Expiration Date: 10/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA

, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0297S/L v1.0

RNase H Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0297SVIAL	RNase H	10123429	Pass	
B0297SVIAL	RNase H Reaction Buffer	10140272	Pass	

Assay Name/Specification	Lot # 10149191
Exonuclease Activity (Radioactivity Release, Single Stranded)	Pass
A 50 µl reaction in RNase H Reaction Buffer containing 1 µg of single stranded [3H]	
E. coli DNA and a minimum of 50 units of RNase H incubated for 30 minutes at 37°C	
releases <0.1 of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in RNase H Reaction Buffer containing 1 µg of supercoiled PhiX174	
DNA and a minimum of 50 units of RNase H incubated for 4 hours at 37°C results in	
<10% conversion to the nicked form as determined by agarose gel electrophoresis.	
qPCR DNA Contamination (E. coli Genomic)	Pass
A minimum of 5 units of RNase H is screened for the presence of E. coli genomic DNA	
sing SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results	
are quantified using a standard curve generated from purified E. coli genomic DNA.	
The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	
Protein Purity Assay (SDS-PAGE)	Pass



M0297S / Lot: 10149191

Page 1 of 2

Assay Name/Specification	Lot # 10149191
RNase H is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of RNase H is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Timothy Meixsell Production Scientist 15 Apr 2022 Michael Tonello

Packaging Quality Control Inspector

15 Apr 2022



M0297S / Lot: 10149191

Page 2 of 2