

www.neb.com info@neb.com



New England Biolabs Certificate of Analysis

Product Name: RNase H
Catalog Number: M0297S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to produce 1

nmol of ribonucleotides from 20 picomoles of a fluorescently labeled 50 base pair RNA-DNA hybrid in a total reaction volume of 50 μ l in

20 minutes at 37°C.

Packaging Lot Number: 10232863
Expiration Date: 12/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA

, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0297S/L v1.0

RNase H Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0297SVIAL	RNase H	10221759	Pass	
B0297SVIAL	RNase H Reaction Buffer	10218041	Pass	

Assay Name/Specification	Lot # 10232863 Pass
Endonuclease Activity (Nicking)	
A 50 µl reaction in RNase H Reaction Buffer containing 1 µg of supercoiled PhiX174	
DNA and a minimum of 50 units of RNase H incubated for 4 hours at 37°C results in	
<10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release, Single Stranded)	Pass
A 50 µl reaction in RNase H Reaction Buffer containing 1 µg of single stranded [³H]	
E. coli DNA and a minimum of 50 units of RNase H incubated for 30 minutes at 37°C	
releases <0.1 of the total radioactivity.	
Protein Purity Assay (SDS-PAGE)	Pass
RNase H is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	
detection.	
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	rass
and a minimum of 1 µl of RNase H is incubated at 37°C. After incubation for 16	



M0297S / Lot: 10232863



Assay Name/Specification	Lot # 10232863
hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of RNase H is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist

15 Dec 2023

Michael Tonello

Packaging Quality Control Inspector

20 Mar 2024



M0297S / Lot: 10232863

Page 2 of 2