

New England Biolabs Certificate of Analysis

Product Name: ProtoScript[®] II Reverse Transcriptase
Catalog Number: M0368L
Concentration: 200,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.
Packaging Lot Number: 10029219
Expiration Date: 08/2020
Storage Temperature: -20°C
Storage Conditions: 20 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.01 % IGEPAL[®] CA-630, 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-M0368S/L/X v1.0

ProtoScript [®] II Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0368LVIAL	ProtoScript [®] II Reverse Transcriptase	10017388	Pass
B1034AVIAL	10X DTT	10019475	Pass
B0368SVIAL	ProtoScript [®] II Reverse Transcriptase Reaction Buffer	0101803	Pass


Assay Name/Specification	Lot # 10029219
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ProtoScript[®] II Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ProtoScript[®] II Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10029219
<p>Protein Purity Assay (SDS-PAGE) ProtoScript® II Reverse Transcriptase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 200 units of ProtoScript® II Reverse Transcriptase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in ProtoScript® II Reverse Transcriptase Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of ProtoScript® II Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Christie Vazquez
Production Scientist
05 Sep 2018



Michael Tonello
Packaging Quality Control Inspector
28 Nov 2018