

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: T4 RNA Ligase 1 (ssRNA Ligase)

Catalog Number: M0437M
Concentration: 30,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to convert 1

nanomole of 5'-[32P] rA16 into a phosphatase-resistant##form in 30

minutes at 37°C.

Packaging Lot Number: 10075389
Expiration Date: 12/2021
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol,

(pH 7.4 @ 25°C)

Specification Version: PS-M0437M v1.0

T4 RNA Ligase 1 (ssRNA Ligase) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
N0437AVIAL	ATP	10061043	Pass	
M0437MVIAL	T4 RNA Ligase 1 (ssRNA Ligase), High Concentration	10061041	Pass	
B1004SVIAL	PEG 8000	10076162	Pass	
B0216SVIAL	T4 RNA Ligase Reaction Buffer	10069909	Pass	

Assay Name/Specification	Lot # 10075389
Endonuclease Activity (Nicking) A 50 μl reaction in T4 RNA Ligase 1 Reaction Buffer containing 1 μg of supercoiled	Pass
PhiX174 DNA and a minimum of 30 units of T4 RNA Ligase 1 (ssRNA Ligase), High Concentration incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in T4 RNA Ligase 1 Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 30 units of T4 RNA Ligase 1 (ssRNA Ligase), High Concentration incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Protein Purity Assay (SDS-PAGE)  T4 RNA Ligase 1 (ssRNA Ligase), High Concentration is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



M0437M / Lot: 10075389

Page 1 of 2

Assay Name/Specification	Lot # 10075389
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of T4 RNA Ligase 1 (ssRNA Ligase), High Concentration is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of RNA transcript and a minimum of 30 units of T4 RNA Ligase 1 (ssRNA Ligase), High Concentration is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Bhairavi Jani

Production Scientist

10 Jul 2020

Michael Tonello

Packaging Quality Control Inspector

10 Jul 2020



M0437M / Lot: 10075389

Page 2 of 2