

New England Biolabs Certificate of Analysis

 Product Name:
 7-deaza-dGTP

 Catalog #:
 N0445S/L

 Concentration:
 5 mM

 Unit Definition:
 N/A

 Lot #:
 0471708

 Assay Date:
 08/2017

 Expiration Date:
 08/2019

Storage Temp:

Storage Conditions: Supplied in Ultrapure water as a lithium salt, (pH 7.0)

-20°C

Specification Version: PS-N0445S/L v1.0 Effective Date: 18 Aug 2017

Assay Name/Specification (minimum release criteria)	Lot #0471708
Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 20 μ l of 7-deaza-dGTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 μ l of 7-deaza-dGTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda DNA, 7-deaza) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2 kb Lambda DNA, 7-deaza) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 2 kb product.	Pass
PCR Amplification (5 kb Lambda DNA, 7-deaza) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM p -Nitrophenyl Phosphate (pNPP) and a minimum of 80 μ l 7-deaza-dGTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0471708
Physical Purity (HPLC) - 7-deaza-dGTP is ≥ 95% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of 7-deaza-dGTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

M.W. Southworth

Authorized by Maurice Southworth 18 Aug 2017







Inspected by
Tony Spear-Alfonso
23 Jan 2018