

## New England Biolabs Certificate of Analysis

**Product Name:** *HindIII*  
**Catalog #:** *R0104T/M*  
**Concentration:** *100,000 units/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Lot #:** *0701509*  
**Assay Date:** *09/2015*  
**Expiration Date:** *9/2017*  
**Storage Temp:** *-20°C*  
**Storage Conditions:** *300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA*  
**Specification Version:** *PS-R0104T/M v1.0*  
**Effective Date:** *29 Oct 2013*

Assay Name/Specification (minimum release criteria)	Lot #0701509
<b>Blue-White Screening (Terminal Integrity)</b> - A sample of Litmus28i vector linearized with a 10-fold excess of HindIII, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.	<b>Pass</b>
<b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of HindIII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 200 units of HindIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 200-fold over-digestion of Lambda DNA with HindIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HindIII.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 60 Units of HindIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> - HindIII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	<b>Pass</b>



---

## New England Biolabs Certificate of Analysis

*\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.*



---

Authorized by  
Derek Robinson  
29 Oct 2013



---

Inspected by  
Katrina Francescone  
30 Sep 2015

