

New England Biolabs Certificate of Analysis


Product Name: AatII
Catalog Number: R0117S
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot Number: 10044764
Expiration Date: 05/2021
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA
Specification Version: PS-R0117S/L v1.0

AatII Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0117SVIAL	AatII	10044765	Pass
B7204SVIAL	CutSmart® Buffer	10042965	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10038711	Pass

Assay Name/Specification	Lot # 10044764
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 60 Units of AatII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of AatII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled Litmus38i DNA and a minimum of 20 Units of AatII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 60 units of AatII incubated for 4	Pass

Assay Name/Specification	Lot # 10044764
<p>hours at 37°C releases <0.1% of the total radioactivity.</p> <p>Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with AatII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AatII.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.



Loren Duquette
Production Scientist
12 Apr 2019



Michael Tonello
Packaging Quality Control Inspector
20 May 2019