

New England Biolabs Certificate of Analysis

Product Name: TaqI-v2
Catalog Number: R0149L
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 65°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10180363
Expiration Date: 02/2025
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 µg/ml BSA, (pH 7.4 @ 25°C)
Specification Version: PS-R0149S/L v2.0


TaqI-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0149LVIAL	TaqI-v2	10180362	Pass
B6004SVIAL	rCutSmart™ Buffer	10179339	Pass

Assay Name/Specification	Lot # 10180363
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of TaqI-v2 incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of TaqI-v2 incubated for 15 minutes at 65°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with TaqI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with TaqI-v2.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 200 units of TaqI-v2 incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10180363
<p>Protein Purity Assay (SDS-PAGE) TaqI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of TaqI-v2 is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
14 Feb 2023



Michael Tonello
Packaging Quality Control Inspector
16 Mar 2023