

## New England Biolabs Certificate of Analysis

**Product Name:** TaqI-v2  
**Catalog Number:** R0149L  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 65°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10211284  
**Expiration Date:** 10/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 µg/ml rAlbumin, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0149S/L v3.0


TaqI-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0149LVIAL	TaqI-v2	10211236	Pass
B6004SVIAL	rCutSmart™ Buffer	10207415	Pass

Assay Name/Specification	Lot # 10211284
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of TaqI-v2 incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of TaqI-v2 incubated for 15 minutes at 65°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with TaqI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with TaqI-v2.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 200 units of TaqI-v2 incubated for 16 hours at 65°C results in a DNA pattern free of	Pass

Assay Name/Specification	Lot # 10211284
<p>detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	
<p><b>Protein Purity Assay (SDS-PAGE)</b> TaqI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of TaqI-v2 is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of TaqI-v2 is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
21 Oct 2023



Michael Tonello  
Packaging Quality Control Inspector  
13 Nov 2023