

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Hgal
Catalog Number: R0154S
Concentration: 2,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of PhiX174 DNA in 1 hour at 37°C in a total reaction volume of 50

μl.

Packaging Lot Number: 10101978
Expiration Date: 03/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0154S/L v1.0

| Hgal Component List    |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0154SVIAL             | Hgal                  | 10101580   | Pass                 |  |
| B7201SVIAL             | NEBuffer™ 1.1         | 10096306   | Pass                 |  |

| Assay Name/Specification   | Lot # 10101978 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE)  | Pass           |
| Hgal is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.  |                |
| Non-Specific DNase Activity (16 Hour)  | Pass           |
| A 50 μl reaction in NEBuffer 1.1 containing 1 μg of PhiX174 DNA and a minimum of 2 Units of Hgal incubated for 16 hours at 37°C results in a DNA pattern free of |                |
| detectable nuclease degradation as determined by agarose gel electrophoresis.  |                |
| Exonuclease Activity (Radioactivity Release)   | Pass           |
| A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and  |                |
| double-stranded [ ³H] E. coli DNA and a minimum of 2 units of Hgal incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.                      |                |
| Ligation and Recutting (Terminal Integrity)  | Pass           |
| After a 2-fold over-digestion of PhiX174 DNA with Hgal, >95% of the DNA fragments  |                |
| can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Hgal.  |                |



R0154S / Lot: 10101978

Page 1 of 2



This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

08 Mar 2021

Michael Tonello

Packaging Quality Control Inspector

08 Mar 2021