

## New England Biolabs Certificate of Analysis

*Product Name:* AflIII  
*Catalog #:* R0541S/L  
*Concentration:* 10,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
*Lot #:* 0321407  
*Assay Date:* 07/2014  
*Expiration Date:* 07/2016  
*Storage Temp:* -20 °C  
*Storage Conditions:* 500 mM NaCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 200 µg/ml BSA  
*Specification Version:* PS-R0541S/L v1.0  
*Effective Date:* 12 Apr 2013

| Assay Name/Specification (minimum release criteria)   | Lot #0321407 |
|---|--------------|
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of AflIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | <b>Pass</b>  |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 5-fold over-digestion of Lambda DNA with AflIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AflIII.   | <b>Pass</b>  |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 30 Units of AflIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.     | <b>Pass</b>  |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

M. W. Southworth

Kerri Sakowski

Authorized by  
Maurice Southworth  
12 Apr 2013

Inspected by  
Kerri Sakowski  
30 Jun 2014

