

## New England Biolabs Certificate of Analysis

**Product Name:** *Nb.BbvCI*  
**Catalog Number:** *R0631S*  
**Concentration:** *10,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to convert 1 µg of supercoiled pUB DNA to open circular form in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Lot Number:** *10044601*  
**Expiration Date:** *05/2021*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*  
**Specification Version:** *PS-R0631S/L v2.0*


Nb.BbvCI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0631SVIAL	Nb.BbvCI	10044604	Pass
B7204SVIAL	CutSmart® Buffer	10042966	Pass

Assay Name/Specification	Lot # 10044601
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 30 units of Nb.BbvCI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Non-Specific DNase Activity (16 hour)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pUB DNA and a minimum of 10 units of Nb.BbvCI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Derek Robinson  
Production Scientist  
08 May 2019



Jay Minichiello  
Packaging Quality Control Inspector  
29 May 2019