

## New England Biolabs Certificate of Analysis

*Product Name:* PI-SceI  
*Catalog #:* R0696S/L  
*Concentration:* 5,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to cleave 1 µg of pBSvdeX XmnI-linearized Control Plasmid in 3 hours at 37°C in a total reaction volume of 50 µl.  
*Lot #:* 0131703  
*Assay Date:* 03/2017  
*Expiration Date:* 3/2019  
*Storage Temp:* -20°C  
*Storage Conditions:* 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA  
*Specification Version:* PS-R0696S/L v2.0  
*Effective Date:* 26 Nov 2013

Assay Name/Specification (minimum release criteria)	Lot #0131703
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in NEBuffer PI-SceI containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of PI-SceI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of pBSvdeX-XmnI DNA with PI-SceI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with PI-SceI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in NEBuffer PI-SceI containing 1 µg of pBSvdeX-XmnI DNA and a minimum of 50 Units of PI-SceI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by  
Derek Robinson  
26 Nov 2013



Inspected by  
Jianying Luo  
17 Mar 2017

