

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Bsal-HF®v2
Catalog Number:	R3733S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10100985
Expiration Date:	03/2023
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 μg/ml BSA, 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-R3733S/L v1.0

Bsal-HF®v2 Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3733SVIAL	Bsal-HF®v2	10100710	Pass	
B7204SVIAL	CutSmart® Buffer	10097266	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10091035	Pass	

Assay Name/Specification	Lot # 10100985
Protein Purity Assay (SDS-PAGE) Bsal-HF®v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 μl reaction in CutSmart® Buffer containing 1 μg of pXba DNA and a minimum of 60 units of Bsal-HF®v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 $\mu$ I reaction in CutSmart® Buffer containing 1 $\mu$ g of supercoiled PhiX174 DNA and a minimum of 20 units of BsaI-HF®v2 incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Bsal-HF®v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10100985
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with Bsal-HF®v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass
fragments can be ligated with 14 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Bsal-HF®v2.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Peph 2 Log

Penghua Zhang Production Scientist 16 Mar 2021

Josh Hersey

Packaging Quality Control Inspector 16 Mar 2021

