240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

Date

11 Jul 2016

www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: AMV Reverse Transcriptase

Catalog #: M0277T

Concentration: 25,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to incorporate 1 nmol of dTTP into an acid-insoluble form in 10

minutes at 37°C.

Shelf Life: 24 months Storage Temp: -20°C

Storage Conditions: 200 mM KPO₄, 2 mM DTT, 0.2 % Triton®X-100, 50 % Glycerol, (pH 7.2 @ 25°C)

Specification Version: PS-M0277T v1.0 Effective Date: 11 Jul 2016

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 µl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 μg of a mixture of single and double-stranded [3 H] E. coli DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 25 units of AMV Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

RNAse Activity Assay (4 Hour Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of AMV Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Derek Robinson

Director of Quality Control





