

## New England Biolabs Product Specification

<b>Product Name:</b>	<i>HindIII</i>
<b>Catalog #:</b>	<i>R0104T/M</i>
<b>Concentration:</b>	<i>100,000 units/ml</i>
<b>Unit Definition:</b>	<i>One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r2.1 in 1 hour at 37°C in a total reaction volume of 50 µl.</i>
<b>Shelf Life:</b>	<i>24 months</i>
<b>Storage Temp:</b>	<i>-20°C</i>
<b>Storage Conditions:</b>	<i>10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)</i>
<b>Specification Version:</b>	<i>PS-R0104T/M v2.0</i>
<b>Effective Date:</b>	<i>08 Jul 2022</i>

### Assay Name/Specification (minimum release criteria)

**Blue-White Screening (Terminal Integrity)** - A sample of Litmus28i vector linearized with a 10-fold excess of HindIII, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

**Endonuclease Activity (Nicking)** - A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of HindIII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 200 units of HindIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Functional Testing (15 minute Digest)** - A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 µl of HindIII incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

**Ligation and Recutting (Terminal Integrity)** - After a 200-fold over-digestion of Lambda DNA with HindIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HindIII.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in NEBuffer™ r2.1 containing 1 µg of Lambda DNA and a minimum of 60 units of HindIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Protein Purity Assay (SDS-PAGE)** - HindIII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.



---

## New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)
---

qPCR DNA Contamination ( <i>E. coli</i> Genomic) - A minimum of 20 units of HindIII is screened for the presence of <i>E. coli</i> genomic DNA using SYBR <sup>®</sup> Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is $\leq 1$ <i>E. coli</i> genome.
--

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Date 08 Jul 2022

---

Derek Robinson  
Director, Quality Control

