

New England Biolabs Product Specification

<i>Product Name:</i>	<i>PI-SceI</i>
<i>Catalog #:</i>	<i>R0696S/L</i>
<i>Concentration:</i>	<i>5,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to cleave 1 µg of pBSvdeX XmnI-linearized Control Plasmid in 3 hours at 37°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA</i>
<i>Specification Version:</i>	<i>PS-R0696S/L v2.0</i>
<i>Effective Date:</i>	<i>26 Nov 2013</i>

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer PI-SceI containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 50 units of PI-SceI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of pBSvdeX-XmnI DNA with PI-SceI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with PI-SceI.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer PI-SceI containing 1 µg of pBSvdeX-XmnI DNA and a minimum of 50 Units of PI-SceI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 26 Nov 2013

Derek Robinson
Quality Approver

