

New England Biolabs Product Specification

<i>Product Name:</i>	<i>PstI-HF[®]</i>
<i>Catalog #:</i>	<i>R3140S/L</i>
<i>Concentration:</i>	<i>20,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton X-100, 200 µg/ml BSA, (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-R3140S/L v2.0</i>
<i>Effective Date:</i>	<i>30 Apr 2018</i>

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart[®] Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 200 units of PstI-HF[®] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 100-fold over-digestion of Lambda DNA with PstI-HF[®], >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PstI-HF[®].

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart[®] Buffer containing 1 µg of Lambda DNA and a minimum of 200 units of PstI-HF[®] incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - PstI-HF[®] is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



Date 30 Apr 2018

Derek Robinson
Director of Quality Control

